Jul

Fill as AD-701077

# The Stability of Coupled Renewal-Differential Equations with Econometric Applications

By

Ronald P. Rhoten and J. K. Aggarwal Department of Electrical Engineering



Technical Report No. 69 July 15, 1969

## INFORMATION SYSTEMS RESEARCH LABORATORY

N72-13508

(NASA-CR-124712) THE STABILITY OF COUPLED

RENEWAL-DIFFERENTIAL EQUATIONS WITH ECONOMETRIC APPLICATIONS R.P. Rhoten, et al (Texas Univ.) 15 Jul. 1969 56 p

Unclas 10234

CSCL 12A

G3/19

The University of Texas at Austin

Austin. Texas 78712

The Electronics Research Center at The University of Texas at Austin constitutes interdisciplinary laboratories in which graduate faculty members and graduate candidates from numerous academic disciplines conduct research.

Research conducted for this technical report was supported in part by the Department of Defense's JOINT SERVICES ELECTRONICS PROGRAM (U.S. Army, U.S. Navy, and the U.S. Air Force) through the Research Grant AF-AFOSR-766-67, Mod. 1 AFOSR-67-0766E. This program is monitored by the Department of Defense's JSEP Technical Advisory Committee consisting of representatives from the U.S. Army Electronics Command, U.S. Army Research Office, Office of Naval Research, and the U.S. Air Force Office of Scientific Research.

Additional support of specific projects by other Federal Agencies, Foundations, and The University of Texas at Austin is acknowledged in footnotes to the appropriate sections.

Reproduction in whole or in part is permitted for any purpose of the U.S. Government.

# THE STABILITY OF COUPLED RENEWALDIFFERENTIAL EQUATIONS WITH ECONOMETRIC APPLICATIONS

Вy

Ronald P. Rhoten and J. K. Aggarwal Department of Electrical Engineering

Technical Report No. 69 July 15, 1969

#### INFORMATION SYSTEMS RESEARCH LABORATORY

ELECTRONICS RESEARCH CENTER
THE UNIVERSITY OF TEXAS AT AUSTIN
Austin, Texas 78712

<sup>\*</sup>Research sponsored in part by the Joint Services Electronics Program under the Research Grant AF-AFOSR-766-67, NSF Grant GK-1879, and NASA Contract NAS8-18120.

# PRECEDING PAGE BLANK NOT FILMED

#### ABSTRACT

This work presents concepts and results in the fields of mathematical modelling, economics and stability analysis. A coupled renewal-differential equation structure is presented as a modelling form for systems possessing hereditary characteristics, and this structure is applied to a model of the Austrian theory of business cycles. For realistic conditions, the system is shown to have an infinite number of poles, and conditions are presented which are both necessary and sufficient for all poles to lie strictly in the left half plane.

## TABLE OF CONTENTS

CHA	PTER																			PAGE
	ABSTRACT .						•	•			•			•			•			iii
	LIST OF FIG	URES	•	•	•		•	•	•		•	•	•	•	•	•	•	•	•	v
I.	INTRODUCT	ION	•	•					•				•	•			•			1
II.	EXISTENCE,	UNIQ	UE	NE	SS	AN	ID	NU	JM	ERI	CA	LI	RES	UI	TS					9
III.	STABILITY C	ONDIT	'IO	NS		•		•				•		•		•				13
IV.	CHARACTERI	STICS	OI	M	IAI	HE	M	ATI	CA	LI	МC	DE	LS	SOI	U	CIO	NS	; .		31
٧.	A ZERO LOCA	ATION	PR	ОВ	LE	M											•	•		44
VI.	CONCLUSIO	NS.								•					•			•		50
BTBT.	IOGRAPHY																			51

### LIST OF FIGURES

FIGUR	LE Company of the Com	PAGE
1	The Numerical Solution	11
2	The Continuing Boom	32
3	The Capital-goods Fluctuation	33
4	The Clyclical Solution	34
5	A Constant Solution	36
6	Index of Value of Building Permits	38
. 7	Department Store Sales	39
8	Government Credit Expansion	40
9	Value of Building Permits Deviation from the Trend	41
10	Department Store Sales Deviation from the Trend	42
11	The Zeros of F(y)	45
12	The Zeros of $G(y)$	47

#### I. INTRODUCTION

### The Time Delay Structure

In order to describe or predict the behavior of a physical system, a common procedure is to represent the various physical variable values at any time t as a finite dimensional vector  $\mathbf{x}(t)$  and to assume that the rate of change of  $\mathbf{x}(t)$  depends only on quantities measurable at time t, leading to the ordinary differential equation system

$$\frac{\mathrm{d}\mathbf{x}(t)}{\mathrm{d}t} \stackrel{\triangle}{=} \mathbf{x}'(t) = f(\mathbf{x}(t), t), \ \mathbf{x}(0) = \mathbf{x}_0. \tag{1.1}$$

The class of physical systems which can adequately be described by (1.1) is large, yet systems in physiology, economics or sociology, among others, often require a more sophisticated mathematical structure. For example, the number of human births at any time clearly depends on the number of women of childbearing age alive nine months previously. Systems of the above form lead to a modelling structure of equations with time delay terms and which may be algebraic, differential or integral in form with single or multiple delays.

There are many examples in the literature of the use of such time delay equations in modelling diverse problems. Lotka [1] considered the problem of industrial replacement with the model

$$p(t) + \int_{0}^{t} f(s) p(t - s) ds = 1,$$
 (1.2)

where p(t) is the probability that an individual machine survives at least tyears, f(t) is the replacement rate and the number of machines in use is constrained to being a constant. Wangersky and Cunningham [2] analyzed the effects of reaction time delay on prey-predator relationships by using as a model the equations

$$x'(t) = A x(t) - A x^{2}(t) / K_{x} - K_{1}x(t) y(t)$$
 (1.3)

$$y'(t) = K_2 \times (t - \tau) y (t - \tau) - my(t),$$
 (1.4)

where x(t) and y(t) are respectively the number of prey and predator at time t,  $\tau$  is the reaction time delay and A,  $K_x$ ,  $K_1$ ,  $K_2$  and m are various constants. A more general form of (1.2), the renewal equation, was used by Lotka [3] in his fundamental work on the dynamic behavior of the age structure of a general population. Denoting by b(t) the number of births at any time t, by b\*(t) the number of births due to the population alive at time zero and by m(t) the maternity function, "Lotka's Equation" is given by

$$b(t) = b^*(t) + \int_0^t b(t - \tau) m(\tau) d\tau.$$
 (1.5)

The structure of (1.5) seems to be especially broad in its applications, having been used for studies in solid state physics [4], plasma dynamics [5] and biology [6]. Such a (vector) form should be considered in any attempt to model large economic processes, since many certainly possess a preponderant hereditary nature. Unfortunately, the analysis of renewal systems becomes increasingly difficult as the system order increases. One purpose of this work is to illustrate how certain simplifying assumptions can be made about a system to reduce this complexity. Specifically, the equation structure to be examined is of the form

$$\underline{x}(t) = \underline{g}(t) + \int_{0}^{t} B(t - \tau) \underline{x}(\tau) d\tau + C\underline{v}(t)$$
 (1.6)

$$\underline{\mathbf{v}}'(t) = \mathbf{D}\underline{\mathbf{x}}(t) + \mathbf{E}\underline{\mathbf{v}}(t), \ \underline{\mathbf{v}}(0) = \underline{\mathbf{v}}_{0}, \tag{1.7}$$

where  $\underline{x}(t)$  and  $\underline{g}(t)$  are n-vectors,  $\underline{v}(t)$  is an m-vector and  $\underline{B}(t)$ ,  $\underline{C}$ ,  $\underline{D}$  and  $\underline{E}$  are  $n \times n$ ,  $m \times n$ ,  $n \times m$  and  $m \times m$  matrices respectively. While at first glance equations (1.6) and (1.7) seem more complex than an ordinary vector renewal equation, they should be considered conceptually as an n order renewal system coupled to an m order differential system to model an m order renewal process. The following section gives the motivation for an economic problem to which the above structure will be applied.

#### Motivation

The business cycle is unarguably a physical process of paramount importance. A "run-away" inflation in Germany at the close of World War I

effectively wiped out the life savings of millions of Germans and the Great Depression caused unbelievable world-wide social agony. Whether the evils of depression exceed those of inflation is a question of at most academic interest; the problem is to control a country's economic behavior between these two extremes. As governments' powers and influence have grown, their abilities to influence a country's economic posture have increased, and the monetary control procedures are now hopefully available. The problem, then, is to recognize at an early time the advent of inflationary or deflationary factors. As an example that such recognition is not at all simple, exactly one week before the crash of 1929, eminent Yale economist Irving Fisher announced that the American economy was moving on a "... permanently high plateau" [7].

Modern economists apply one of two methods in their "prognostic" attempts. The first of these involves the use of econometric models of an empirical-statistical nature. All such models have in common a relatively short-term forecastive ability, which may or may not be a liability, and an almost overwhelming complexity, certainly a liability. The Wharton Econometric and Forecasting Unit model, for example, consists of 52 stochastic equations, 29 identities, 144 statistically determined variables and 117 endogenous and exogenous variables [8]. Such complexity, besides causing computational problems, also tends to obscure the possible existence of fundamental economic factors which, if completely understood, might lead to a more basic understanding of the economic process. These substrative concepts form the basis of the second forecastive approach, a more qualitative treatment.

Qualitative economic theories attempt to present a foundation that simply describes the system's aggregate behavior, yet is comprehensive enough to allow explanation of the myriad of peripheral activities. Since Adam Smith's Wealth of Nations [9], countless theories have been propounded,

and some, like Smith's, are completely valid given the circumstances under which they were formulated. Changing conditions, however, have shown even the best of these to be dated. To discover whether or not any particular theory will prove independent of the events surrounding its inception, it should be modelled mathematically. This allows a testing procedure far more general than a mere comparison with past history. Although some attempts have been made along these lines (see in particular reference [8], Chapter 14), there still exist many qualitative theories which have not been quantitatively expressed. This work will present a simple model of one of these, the Austrian theory [10] of business cycles.

The Austrian theory, as expounded by Ludwig von Mises [11] and F. A. Hayek [12], is basically a monetary theory of the cycle. If an economy has a fixed supply of money, that amount available for investment is a function of the populace's time preference for consumption. If the people are in a buying mood, only a small percentage of their income is "saved" through savings accounts, insurance policies, stocks and bonds, and the interest rate will rise. If, on the other hand, the time preference is slanted toward future consumption, larger savings are made and the interest rate drops. Turning now to the other half of the investment process, the Austrian theory divides industry into two sections, the capital goods industries and the consumer-goods industries. The capital-goods section is characterized mainly by its long-term aspects. The building of a steel mill, for example, requires considerable time and continued investments as work progresses. Such investments, then, should be made at low interest rate conditions, since these indicate future consumer desire. The consumer-goods industries, conversely, are more quickly expanded, and should expand when the interest rate is high, indicating an immediate demand for goods. The overall process seems stable, since the availability of consumer goods is increased when the demand for them is high, and the entrepreneur builds for future consumption when society is saving.

Suppose now that the government intrudes into this investment process by credit expansion through federal banks. As credit is created, the interest rate drops and it appears to the entrepreneur that consumer time preferences have lengthened. This clearly calls for a shifting to capital-goods expansion, but as the excess money filters down to the consumer, he attempts to assert this increase in buying power in accordance with his old saving/ consumption proportions. This, of course, completely startles the business community, which had been misled by the drop in interest rates into believing that these proportions had changed. As soon as the government decides to end its inflationary credit policies, as it must sooner or later, the continued investments required for the capital-goods industries are not obtainable at profitable interest rates, and previous investments are found to have been illconceived. The only recourse is liquidation of certain projects, signalling the beginning of the depression. The recovery period begins only when the capital-goods/consumer-goods investment proportions are in agreement with the consumption characteristics of the individual buyer.

While this is only a brief summary of the Austrian theory, it does cover the main concepts and provides a starting point for the mathematical modelling process.

# Formulation of Model and Summary of Results

The model to be presented will characterize the Austrian theory at least locally. That is, a linear model will be formulated in an attempt to describe small deviations of the physical process from the trend. Suppose  $\mathbf{x}(t)$  is defined to be the investment in capital-goods industries differing from the trend, and consider the viewpoint the entrepreneur takes in deciding whether or not to invest in  $\mathbf{x}(t)$ . Recalling that such industries have long completion times, our entrepreneur would certainly consider the activity in this sector over the past several years. The existence of several steel mills under construction would, of course, tend to dampen interest in this area. This could be expressed mathematically by

$$x(t) \propto -\int_0^t \rho(\tau) x (t - \tau) d\tau, \qquad (1.8)$$

where  $\rho(t)$ , non-negative in general, might be called an "influence" function. If i(t) is defined as the difference between the current rate of interest and the trend, the theory states that x(t) will tend to be positive when i(t) is negative; thus,

$$x(t) \propto -i(t). \tag{1.9}$$

Turning now to the investment in consumer-goods industries as differing from the trend, denoted by v(t), it would be possible to structure a modelling equation exactly as (1.8) and (1.9). However, the time-delays in such industries are of comparatively short duration, and the system model is simplified if it is assumed that this faster response process can be modelled by an ordinary differential equation. That is, since an investor in a consumer-goods industry should change his position with respect to the instantaneous levels of investment activity, the model should express

$$v'(t) \propto -v(t). \tag{1.10}$$

This investment behavior as a function of the interest rate is also of a somewhat different character. Since consumer-goods production can increase so rapidly, a high interest rate should signal further increases; that is, the derivative of v(t) should be proportional to i(t),

$$v'(t) \propto i(t)$$
. (1.11)

While the system of proportionalities (1.8) - (1.11) could now be combined in a mathematical model, a more appropriate system of equations results from an additional consideration. The changes are to be concerned with i(t), since the most modern economic theories are not so directly tied to the interest rate "per se". There is certainly a question of the entrepreneur's ability to recognize the interest rate trend, and thus, use deviations from this trend in any decision process. Perhaps the best approximation to the

business community's behavior is to assume that information concerning the interest rate is actually gained through a study of the existing market investment structure. That is, increases in capital-goods investment are tied more directly to decreases in consumer-goods investment than to any interest rate deviations. This line of reasoning leads to a change in proportionalities (1.9) and (1.11) to

$$x(t) \propto -v(t) \tag{1.12}$$

and

$$v'(t) \propto -x(t)$$
. (1.13)

The government's credit expansion policies may now be denoted separately by g(t) and considered as a system input,

$$x(t) \propto g(t) \tag{1.14}$$

Combining (1.8), (1.10), (1.12), (1.13) and (1.14) with appropriate constants of proportionality, there results

$$x(t) = g(t) - \int_{0}^{t} \rho(t - \tau) x(\tau) d\tau - \alpha v(t)$$
 (1.15)

$$v'(t) = -\gamma x(t) - \beta v(t), v(0) = v_0.$$
 (1.16)

These equations are structurally identical to (1.6) and (1.7), and the modelling process has indicated the reduction to a minimum of the order of the renewal equation portion of the mathematical model.

With the qualitative Austrian theory now quantitatively expressed, the questions become mathematical in nature. In Chapter II, equations (1.6) and (1.7) are imbedded in a more general structure and existing theorems utilized to provide conditions which insure the existence, uniqueness and transformability of solutions. Some discussion is given concerning the calculation of numerical results, with the differential character of (1.6) - (1.7) emphasized. Chapter III is devoted to a system stability analysis. With the determinental equation expressed as a finite exponential series, necessary

and sufficient conditions for all zeros to lie strictly in the left half plane are presented. The results are a generalization of those of Pontryagin [13] for exponential polynomials. Finally, an application is presented to illustrate the use of the stability conditions, and additional economic implications are mentioned.

# II. EXISTENCE, UNIQUENESS AND NUMERICAL RESULTS

This section will present conditions which will insure the existence, uniqueness and transformability of solutions. To utilize existing theorems, system (1.6) - (1.7) will be imbedded in a more general structure. Equation (1.7) may be integrated on both sides and, with the appropriate initial condition, becomes

$$\underline{\mathbf{v}}(t) - \underline{\mathbf{v}}_0 = \int_0^t \left\{ D \underline{\mathbf{x}}(\tau) + \mathbf{E} \underline{\mathbf{v}}(\tau) \right\} d\tau. \tag{2.1}$$

Solving (2.1) for  $\underline{v}(t)$  and substituting the result in (1.6) yields

$$\underline{\mathbf{x}}(t) = \underline{\mathbf{g}}(t) + \mathbf{C} \underline{\mathbf{v}}_0 + \int_0^t \mathbf{B}(t - \tau) \underline{\mathbf{x}}(\tau) d\tau + \mathbf{C} \int_0^t \left\{ \mathbf{D} \underline{\mathbf{x}}(\tau) + \mathbf{E} \underline{\mathbf{v}}(\tau) \right\} d\tau$$

and thus

$$\begin{bmatrix} \underline{\mathbf{x}}(t) \\ \underline{\mathbf{v}}(t) \end{bmatrix} = \begin{bmatrix} \underline{\mathbf{g}}(t) + \mathbf{C}\underline{\mathbf{v}}_0 \\ \underline{\mathbf{v}}_0 \end{bmatrix} + \int_0^t \begin{bmatrix} \mathbf{B}(t-\tau) + \mathbf{C}\mathbf{D} & \mathbf{C}\mathbf{E} \\ \mathbf{D} & \mathbf{E} \end{bmatrix} \begin{bmatrix} \underline{\mathbf{x}}(\tau) \\ \underline{\mathbf{v}}(\tau) \end{bmatrix} d\tau.$$
(2.2)

Hence, equations (1.6) - (1.7) may be expressed in the vector form

$$\underline{z}(t) = \underline{h}(t) + \int_0^t A(t - \tau) \underline{z}(\tau) d\tau. \qquad (2.3)$$

It is of interest to note that the initial condition,  $\underline{v}_0$ , appears in (2.1) in a manner analogous to  $\underline{g}(t)$  in (1.6). Although  $\underline{g}(t)$  is just the input,  $\underline{g}(0) + C\underline{v}_0$  is also the initial value of  $\underline{x}(t)$ . If it is desired to arbitrarily set an initial condition for  $\underline{x}(t)$ ,  $\underline{g}(t)$  must be defined appropriately for t=0 or  $\underline{x}(0)$  will be multiply defined. Thus, the input set for (2.3) also defines the space of applicable initial conditions.

The proofs of the following theorems are extensions to the vector case of available proofs [14] for the scalar form of (2.3), and will not be presented here. Introduce the vector and matrix norms

$$\left\|\underline{z}\right\| = \sum_{i=1}^{n} |z_{i}|$$

and

$$||A|| = \sum_{i,j=1}^{n} |a_{ij}|.$$

Theorem 2.1. If

(i)  $\|\underline{h}(t)\| \le c$  for all  $t \in [0, t_0]$  and

(ii) 
$$\|A(t)\| \le m$$
 for all  $t \in [0, t_0]$ ,

then the solution to (2.3) exists and is unique for t  $_{6}[0,t_{0}]$ , and is continuous if  $\underline{h}(t)$  is continuous.

Theorem 2.2. Let h(t) be continuous. If for some

a there exist two positive constants  $c_1$  and

c, such that

(i)  $\|\underline{h}(t)\| \le c_1 e^{at}$  for all t > 0, and

(ii) 
$$\int_0^\infty e^{-at} ||A(t)|| dt = c_2 < 1$$
,

then

$$\|\underline{z}(t)\| \leq \frac{c_1 e^{at}}{1 - c_2}$$

for all  $t \ge 0$ , and the Laplace transform of  $\underline{z}(t)$  exists.

The basic tool in the proof of the above theorems is the technique of successive approximations. This technique may also be used to find conditions which insure that  $\underline{z}(t)$  is Lesbeque integrable, Riemann integrable or of bounded variation ([14], pp. 217-29).

The foregoing imbedding has emphasized the integral characteristics of equations (1.6) and (1.7). Their differential character is perhaps best illustrated by noting the simplifications in numerical solution techniques which the differential character of (1.7) makes possible. Figure 1 illustrates

#### THE NUMERICAL SOLUTION




Figure 1

the utilization made of a sophisticated, fast differential equation solving subroutine [15] in solving (1.15) and (1.16) for particular values of g(t),  $_{\rho}(t), \,_{\alpha}, \,_{\beta}, \gamma$  and  $v_0$ . For a solution desired for  $0 \leq t \leq T$ , an  $\,$  N is chosen such that  $t_0=0, \,\,t_n=T$  and  $\Delta t,$  defined by  $\Delta t=t_{i+1}-t_i, \,\,i< N,$  is sufficiently small to allow good approximations to (1.15) by a rectangular integration process with rectangles of width  $\Delta t.$  The input and influence function values are stored for appropriate values of i, and an error specification set for the subroutine (DEQB). DEQB is then entered with the initial value  $v_0$  and with i=0, and instructed to find  $v(t_{i+1})$  under the constraints of (1.16). At various times  $t_0$ ,  $t_i < t_0 < t_{i+1}$ , control is returned to the main program to evalutate  $v'(t_0)$ . The input,  $g(t_0)$ , is found by linear interpolation and the convolution integral approximated by

$$\Delta t \stackrel{i}{\sum}_{i=0} x(t_{j}) \rho(t_{i-j+1}) + \Delta t \rho_{0}(x(t_{i}) + (x(t_{i}) - x(t_{i-1}))(t_{0} - t_{i}) / \Delta t).$$

The subroutine itself provides an initial guess for  $v(t_0)$ , so  $v'(t_0)$  is now calculated and control returned to DEQB. As the exit is made from DEQB at  $t_0 = t_{i+1}$ ,  $v(t_{i+1})$  and  $x(t_{i+1})$  are available for output,  $x(t_{i+1})$  is stored, i is incremented and control is returned to the subroutine for the next iteration.

While memory storage needs are large for evaluation of the convolution integral, additional differential equations could be added to (1.15) and (1.16) with only linearly increasing requirements. The subroutine provides automatic step size correction to minimally meet specified error contraints, which leaves only the evaluation of the integral for the programmer. While only a crude rectangular approximation was used for demonstration purposes, even rather critical constant solutions were evaluated within acceptable limits. These and other solutions will be presented later as examples.

#### III. STABILITY CONDITIONS

With conditions now having been given insuring the existence, uniqueness and exponential boundedness of solutions, the main questions remaining concern system stability. The results to be presented are quite general, applying to (1.6) – (1.7), but algebraic considerations will require use of equations (1.15) and (1.16) in illustrative examples.

If g(t) and  $\rho$ (t) in (1.15) - (1.16) satisfy the conditions which allow application of the Laplace Transform, and if  $v_0$  = 0, then

$$\begin{bmatrix} X(z) \\ V(z) \end{bmatrix} = \begin{bmatrix} 1 + P(z) & \alpha \\ \gamma & z+\beta \end{bmatrix} - 1 \begin{bmatrix} G(z) \\ 0 \end{bmatrix} ,$$

and the system stability is clearly a function of the location of the zeros of the determinantal equation.

$$D(z) = z + z P(z) + \beta P(z) + (\beta - \alpha \gamma).$$
 (3.1)

Since the difficulty in finding the zero locations of (3.1) depends on the form of P(z), a particular  $\rho(t)$ ,

$$\rho(t) = \begin{cases} \sum_{j=0}^{m} \rho_{j} t^{j}, & 0 \le t \le T \\ 0, & T < t \end{cases}$$
 (3.2)

will be considered. This form is chosen for two reasons. First, the hereditary character of the system can certainly be considered to extend back in time only some finite length. Second, by the Stone-Weierstrass theorem [16], any function continuous on any closed and bounded interval [a,b] may be approximated uniformly there by a polynomial. Thus,

$$P(z) = \sum_{j=0}^{m} \int_{0}^{T} \rho_{i} t^{i} e^{-zt} dt$$

$$= -\frac{\rho_0 e^{-zT}}{z} + \frac{\rho_0}{z} - \rho_1 \left\{ \frac{T e^{-zT}}{z} + \frac{e^{-zT}}{z^2} \right\} + \frac{\rho_1}{z^2}$$

$$- \dots - \rho_{m} e^{-zT} \int_{i=0}^{m} \frac{i!T}{z^{i+1}} + \frac{\rho_{m}m!}{z^{m+1}}$$

$$= \sum_{j=0}^{m} \left\{ -\rho_{j} e^{-zT} \sum_{i=0}^{j} \frac{i!T^{j-i}}{z^{i+1}} + \frac{\rho_{j}^{j!}}{z^{j+1}} \right\}.$$
 (3.3)

Then

$$D(z) = z + (z + \beta) \left\{ \sum_{j=0}^{m} \left[ -\rho_{j} e^{-zT} \sum_{i=0}^{j} \frac{i!T^{j-i}}{z^{i+1}} + \frac{\rho_{j}j!}{z^{j+1}} \right] \right\} + (\beta - \alpha \gamma).$$
(3.4)

Since the zeros of D(z) are the quantities of interest, define  $\bar{D}(z) = e^{zT}D(z)$  and the zeros of  $\bar{D}(z)$  are just those of D(z). Hence,

$$\bar{D}(z) = \{ z + (\beta - \alpha \gamma) + (z + \beta) \sum_{j=0}^{m} \frac{\rho_j j!}{z^{j+1}} \} e^{zT} 
- (z + \beta) \{ \sum_{j=0}^{m} \rho_j [\sum_{j=0}^{j} \frac{i! T^{j-1}}{z^{j+1}} ] \}.$$
(3.5)

It is easily seen that  $\bar{D}(z)$  may, in general, have an infinite number of zeros. This causes two serious obstacles to arise in stability studies. First, of course, is the difficulty in locating "all" of the zeros. Then, even if it is known that all the zeros have strictly negative real parts, they may approach the imaginary axis asymptotically, giving rise to undamped solutions. Compared to the first problem, this latter one might be considered somewhat pathological, although it will not be neglected in the ensuing work.

Perhaps the most useful results available in the area of zero location for transcendental functions are those due to Pontryagin [13]. Basically, he gives necessary and sufficient conditions for an exponential polynomial to have all of its zeros strictly in the left half plane. Considering (3.5), it is seen that  $\bar{D}(z)$  is not an exponential polynomial, and the obvious procedure would be to examine the zeros of  $z^{m+1}$   $\bar{D}(z)$ . This approach would certainly be valid if the zeros of  $\bar{D}(z)$  and  $z^{m+1}\bar{D}(z)$  were coincident; that is, if  $\bar{D}(z)$  had an  $(m+1)^{st}$  order pole at the origin. A simple counterexample shows, however, that this supposition is not always true. Consider (3.5) for a  $\rho(t)$  of the form  $\rho_n t^n$  for  $0 \le t \le T$  and zero elsewhere. Then an evaluation at the origin yields

$$\begin{split} \bar{D}(z)\big|_{z=0} &= \frac{\text{Lim}}{z^{n+2}} \; \{ \frac{\left[z^{n+2} + (\beta - \alpha \gamma) \; z^{n+1} + n! \, \rho_n z + n! \, \beta \rho_n\right] \, \mathrm{e}^{zT}}{z^{n+1}} \\ &- \frac{(\beta \rho_n + z \rho_n) \, \left(\sum\limits_{i=0}^{n} i! \; T^{n-i} z^{n-i}\right)}{z^{n+1}} \; \} \end{split}$$

which is indeterminate of the form 0/0. Applying L'Hospital's Rule,

$$\begin{split} \bar{D}(z)\big|_{z=0} &= \frac{\text{Lim}}{z \to 0} \; \big\{ \; T \; \big[z^{n+2} + (\beta - \alpha \gamma) \; z^{n+1} + n! \rho_n z + n! \beta \rho_n \big] \, e^{zT} \\ &+ \; \big[ (n+2) \; z^{n+1} + (n+1) \, (\beta - \alpha \gamma) \; z^n + n! \; \rho_n \big] \, e^{zT} \\ &- \; (\beta \rho_n + z \rho_n) \, \sum_{i=0}^{n-1} \, (n-i)i! \; T^{n-i} \; z^{n-i-1} \\ &- \; \rho_n \, \sum_{i=0}^{n} \; i! \; T^{n-i} \; z^{n-i} \; \big\} \; / \, \sum_{z \to 0}^{\text{Lim}} \; \big\{ (n+1) \; z^n \big\} \\ &= \frac{\text{Lim}}{z \to 0} \; \big\{ \frac{[n! \; - \; (n-1) \; ! \; ] \; T \; \beta \rho_n}{(n+1) \; z^n} \big\} \; , \; n \neq 0 \; , \end{split}$$

and  $\bar{D}(z)$  is seen to have at most an  $n^{th}$  order pole at the origin. Thus,

 $z^{n+1}\bar{D}(z)$  will have at least a first order zero at the origin that  $\bar{D}(z)$  does not have. Unfortunately, use of Pontryagin's results will only imply that under no conditions can  $z^{n+1}\bar{D}(z)$  possess only left half plane zeros, and there is no way provided to treat the extraneous zero separately.

Since exponential polynomials have been shown to be of insufficient generality for the study at hand, consideration will now be given to the zero location problem of the next more general category of transcendental functions, the finite exponential series. That is, transcendental functions of the form

$$H(z) = \sum_{m=-q}^{p_1} \sum_{n=0}^{p_2} a_{mn} z^m e^{nz}$$
 (3.6)

will be considered. This set of functions clearly includes the determinental equation of (1.6) - (1.7) for influence functions of the form of (3.2), and in fact is of sufficient generality to allow influence functions sectionally describable as (3.2) and hence only piecewise continuous. Such time sections, however, must be rationally related, since the restriction of n in (3.6) to integer values will require all time intervals to be integer multiples of some base interval. Finite exponential series, as defined by (3.6), are closely related to exponential polynomials, and the following results will be a generalization of Pontryagin's basic work. The proofs of some of his theorems will be utilized directly, others changed and still others corrected.

Two basic lemmas which will be used later will now be proven.

Lemma 3.1. If  $v^{(s)}(t)$  is any non-zero polynomial in t of degree s, then there exists a real number  $\varepsilon$  such that  $v^{(s)}(\exp{(x+i(\varepsilon+2n\pi))})$  is non-zero for all real x and all integer n.

Proof. Since  $v^{(s)}(\exp z)$  is periodic with period  $2\pi i$ , the result will hold for all integer n if it holds for n=0. There clearly exist only s complex numbers  $t_0$  which satisfy

$$v^{(s)}(t_0) = 0.$$

Then  $v^{(s)}(\exp z_0) = 0$  if and only if  $\exp(z_0) = t_0$ . For y(z = x + iy) in the interval  $y_0 \le y \le 2\pi + y_0$ , any  $y_0$ , there are at most two such  $z_0$ 's for each  $t_0$ . Hence,  $v^{(s)}(\exp z)$  has only 2s zeros in any semi-infinite horizontal strip, of width  $2\pi$ , in the complex plane. Thus, there exists an infinity of numbers  $\varepsilon$ , and in particular one, such that  $v^{(s)}(\exp(x + i\varepsilon)) \ne 0$  for all real x.

Q.E.D.

Lemma 3.2. If  $\delta(z)$  is any polynomial in cos z and sin z, of degree s, not identically zero, then there exists an  $\varepsilon$  such that  $\delta(\varepsilon + 2n\pi + iy)$  is non-zero for all real y and integer n.

Proof.  $\delta(z)$  is clearly periodic with period  $2\pi$ , and if the result holds for n=0, it therefore holds for all n. Using a common variable substitution, let

$$u = (t + 1/t) / 2, \quad v = (t - 1/t) / 2i$$
 (3.7)

and for  $t = \exp(iz)$ ,  $u = \cos z$  and  $v = \sin z$ . Then substituting (3.7) for u and v in  $\delta(u,v)$ , the resultant  $\delta(t)$  clearly has 2s zeros. Every zero,  $z_0$ , of  $\delta(z)$  must thus satisfy  $\exp(iz_0) = t_0$ , where  $t_0$  is a zero of  $\delta(t)$ . Hence, for x in the interval  $x_0 \le x \le x_0 + 2\pi$ ,  $\delta(z)$  has at most 4s zeros, and there exists an infinite set of numbers  $\varepsilon$  such that  $\delta(\varepsilon + iy) \ne 0$  for all y. Q.E.D.

Now define h(z,t), a finite series in the two variables z and t, by

$$h(z,t) \stackrel{\Delta}{=} \stackrel{p_1}{\Sigma} \stackrel{p_2}{\Sigma} a_{mn} z^m t^n.$$

The finite exponential series, H(z), may thus be defined  $H(z) \stackrel{\triangle}{=} h(z, \exp(z)). \tag{3.8}$ 

From the discussions of the previous section, it will be assumed without loss of generality that H(z) possesses no pole at the origin. It must also be assumed throughout the remaining discussions that H(z) is not degenerate; i.e., not only the coefficients  $a_m$  are non-zero and there exists an  $m_0 > 0$  such that  $a_m \neq 0$  for some  $n \neq 0$ .

If there exists a non-zero coefficient  $a_{rs}$  such that for all other non-zero  $a_{mn}$ , r>m and  $s\geq n$  or  $r\geq m$  and s>n, then  $a_{rs}$   $z^rt^s$  will be called the principal term of the finite series. This definition leads to the following instability result.

Theorem 3.1. If h(z,t) has no principal term, then H(z) has an infinite number of zeros with arbitrarily large positive real parts.

Proof. Write

$$z^{q}h(z,t) = \sum_{m=q}^{p_1} \sum_{n=0}^{p_2} a_{mn}z^{m+q}t^n;$$

 $z^q$ H(z) is thus an exponential polynomial. If h(z,t) has no principal term, then neither does  $z^q$ h(z,t). By Pontryagin's first theorem,  $z^q$ H(z) thus has an infinite number of zeros with arbitrarily large positive real parts, and H(z) certainly possesses the same property. Q.E.D.

If h(z,t) does possess a principal term, a consideration of the behavior of H(z) along the imaginary axis will yield information concerning the location of the zeros. For z imaginary, H(z) may be separated into its real and imaginary parts by

$$H(iy) = \sum_{m=-q}^{p_1} \sum_{n=0}^{p_2} a_{mn} i^m y^m \exp(iny)$$
(3.9)

$$= \sum_{m=-q}^{p_1} \sum_{n=0}^{p_2} a_{mn} i^m y^m (\cos y + i \sin y)^n.$$
 (3.10)

Then H(iy) may be written

$$H(iy) = \sum_{m=-q}^{p_1} \sum_{n=0}^{p_2} y^m \left[ \phi_m^{(n)}(\cos y, \sin y) + i \psi_m^{(n)}(\cos y, \sin y) \right]$$
(3.11)

where  $\phi_m^{(n)}$  (u,v) and  $\psi_m^{(n)}$  (u,v) are polynomials of degree n, homogeneous in u and v. F(y) and G(y) are now defined by

$$F(y) \stackrel{\Delta}{=} \begin{array}{cccc} p_1 & p_2 & m & p_m \\ \sum & \sum & p_m & p_m \end{array} (cosy, siny), \qquad (3.12)$$

$$m = -q \quad n = 0$$

$$G(y) = \sum_{m=-q}^{\Delta} \sum_{n=0}^{p_2} y^m \psi^{(n)} (\cos y, \sin y), \qquad (3.13)$$

and hence

$$H(iy) = F(y) + i G(y).$$
 (3.14)

Examining the structure of F(y) and G(y), certain observations can be made which will be useful later. F(y) and G(y) will clearly have no pole at the origin. Suppose a function p(z,u,v) is defined by

$$p(z,u,v) \stackrel{\Delta}{=} \stackrel{p_1}{\Sigma} \stackrel{p_2}{\Sigma} \stackrel{z}{\Sigma} \stackrel{z^m}{\gamma_m} (u,v)$$

$$\stackrel{m=-q}{=} n=0$$
(3.15)

where  $\gamma_m^{(n)}(u,v)$  is a polynomial of degree n, homogeneous in u and v, and  $\gamma_m^{(n)}(\cos y, \sin y)$  is formed as  $\phi_m^{(n)}(\cos y, \sin y)$  and  $\psi_m^{(n)}(\cos y, \sin y)$  in (3.13). Then F(y) and G(y) may be expressed as  $f(y, \cos y, \sin y)$  and  $g(y, \cos y, \sin y)$  where f(z,u,v) and g(z,u,v) are of the form of p(z,u,v). Suppose further that a principal term for p(z,u,v) is defined as expected. Then if h(z,t) has a principal term a  $z^r t^s$ , f(z,u,v) and g(z,u,v) will have principal terms  $z^r t^s$  (s)  $z^r t^s$  (u,v) and conversely.

Suppose p(z,u,v) has a principal term  $z \gamma \binom{s}{r} (u,v)$ . Then define

$$\gamma_{\star}^{(s)}(u,v) \stackrel{\Delta}{=} \sum_{r=0}^{s} \gamma_{r}^{(n)}(u,v)$$
 (3.16)

and

$$\frac{-}{Y_{\star}}(s)_{(z)} = \frac{\Delta}{Y_{\star}}(s)_{(\cos z, \sin z)}. \tag{3.17}$$

Such definitions lead to the following fundamental theorem.

Theorem 3.2. Let p(z,u,v) have a principal term  $z \gamma_r^{(s)}(u,v)$ , and choose an  $\varepsilon$  such that  $\gamma_*^{(s)}(\varepsilon + zn\pi + iy)$  is never zero for any real y. Define  $P(z) = p(z, \cos z, \sin z)$  and assume that P(z) has no pole at the origin. Then for k sufficiently large, P(z) will have exactly  $4 \le k + r$  zeros in the strip  $-2k\pi + \varepsilon \le x \le 2k\pi + \varepsilon$ .

Proof. If  $\gamma_{\star}^{-}(s)$  can be shown to be not identically zero for all z. Lemma 2 will imply the existence of an appropriate  $\varepsilon$ . Since  $\gamma_{\star}^{-}(s)$  (z) merely represents either  $\varphi_{\star}^{-}(s)$  (z) or  $\psi_{\star}^{-}(s)$ , it may be shown that neither of these functions is zero for all z. Recalling (3.9), it is clear that an alternate form for (3.10) may be given by

$$H(iy) = \sum_{m=-q}^{p_1} \sum_{n=0}^{p_2} a_{mn} i^m y^m (\cos(ny) + i\sin(ny)).$$
 (3.18)

This implies the following:

if 
$$m = \dots, -4, 0, 4, \dots$$
 then
$$\varphi \begin{pmatrix} (n) \\ m \end{pmatrix} (\cos y, \sin y) = a_{mn} \cos(ny)$$

and

$$\psi_{m}^{(n)}$$
 (cosy, siny) =  $a_{mn}$  sin(ny);

if 
$$m = \dots, -1, 3, 7, \dots$$
 then
$$\varphi \binom{(n)}{m} (\cos y, \sin y) = a_{mn} \sin(ny)$$

and

$$\psi_{m}^{(n)}$$
 (cosy, siny) =  $-a_{mn}$  cos(ny);

if m = ..., -2, 2, 6, ... then

$$\varphi \binom{(n)}{m} (\cos y, \sin y) = -a_{mn} \cos(ny)$$

and

$$\psi_{m}^{(n)}$$
 (cosy, siny) =  $-a_{mn}$  sin(ny);

if m = ..., -3, 1, 5, ... then

$$\varphi \frac{(n)}{m}(\cos y, \sin y) = -a_{mn} \sin(ny)$$

and

$$\psi_{m}^{(n)}$$
 (cosy, siny) =  $a_{mn}$  cos(ny).

Suppose the principal term of H(iy) is  $a_{rs}^{i} y^{r} e^{isy}$ , and for definiteness suppose r = 4k, some k. Then

$$\varphi_{*}^{(s)}(z) = a_{r0}^{+} a_{r1}^{-} \cos z + \dots + a_{rs}^{-} \cos(sz)$$
 (3.19)

For  $\varphi_*^{-(s)}(z)$  to be zero for all z, it clearly must be zero for z = ix. Equation (3.19) then becomes

$$\frac{-(s)}{\varphi_{*}}(z) = a_{r0} + a_{r1} \cosh x + \dots + a_{rs} \cosh(sx)$$
. (3.20)

But since  $\cosh x$  is monotonically increasing for  $x \ge 0$  and  $\cosh n_1 \times \cosh n_2 \times \text{ for } n_1 > n_2 \text{ and } x > 0$ , (3.20) clearly cannot be identically zero. Similar arguments hold for  $\psi_*$  (z) and for r = 4k+1, 4k+2 and 4k+3, any k. With the existence of an appropriate  $\varepsilon$  now shown, and proceeding in a fashion similar to that in the proof of Theorem 3.1, write

$$z^{q}p(z,u,v) = \sum_{m=-q}^{p} \sum_{n=0}^{p} z^{m+q} \gamma^{(n)}(u,v).$$

Then since p(z,u,v) has a principal term  $z \gamma_r^{(s)}(u,v)$ ,  $z_p^{(q)}(z,u,v)$  will have a principal term  $z \gamma_r^{(q)}(u,v)$ .

Pontryagin's third theorem may then be used to imply that  $z^q P(z)$  has, for sufficiently large k, 4sk+r+q zeros in the strip  $-2k\pi+e \le x \le 2k\pi+e$ . Since P(z) has no pole at the origin, it thus has 4sk+r zeros in the same strip, again for sufficiently large k. Q.E.D.

A similar extension of Pontryagin's second theorem leads to the following result.

Theorem 3.3. If p(z,u,v) has no principal term, then P(z) has an infinite number of non-real zeros.

Suppose now that h(z,t) has a principal term  $a_{rs}z^{r}t^{s}$ . Denote by  $\eta_{\star}^{(s)}(t)$  the polynomial coefficient of  $z^{r}$ . Then h(z,t) may be written

$$h(z,t) = z^{r} \eta_{\star}^{(s)}(t) + \sum_{m=-q}^{r-1} \sum_{n=0}^{s} a_{mn} z^{m} t^{n}.$$
 (3.21)

Next, denote by  $N_k$  the number of zeros of H(z) in the semi-infinite strip defined by  $x \ge 0$ ,  $-2k\pi + \varepsilon \le y \le 2k\pi + \varepsilon$ . Denote by  $\omega$  (a,b) the overall angular variation of the vector  $\omega = H(iy)$  about the origin as y ranges from a to b.

Theorem 3.4. Let h(z,t) have a principal term  $a_{rs}s^tt^s$  and choose an  $\varepsilon$  such that  $\eta^{(s)}_*$  (exp(x + i (2n $\pi$  +  $\varepsilon$ ))) is non-zero for all positive, real x and integer n. Assume that H(z) has no zeros on the imaginary axis and no pole at the origin. Then

$$\omega^*(-2k\pi + \epsilon, 2k\pi + \epsilon) = 2\pi (2sk - N_k + r/2) + \delta_k$$
(3.22)

where  $\delta_k \to 0$  as  $k \to \infty$ .

Proof. By the fundamental theorem of algebra,  $\eta_*^{(s)}(t)$  can be identically zero only if it is the zero polynomial. But this would imply that  $a_{rs} = 0$ , a contradiction by

definition of a principal term. By Lemma 3.1 then, an appropriate  $\varepsilon$  may be chosen. Since for z large the leading term of H(z) is dominant, and H(z) has been previously assumed to be non-degenerate, H(z) can be written

$$H(z) = z^{r} \eta_{*}^{(s)} (\exp(z)) [1 + \delta_{1}]$$
 (3.23)

where  $\delta_1 \rightarrow 0$  as  $|z| \rightarrow \infty$ . By Lemma 3.1, there are clearly no zeros of  $z^r \eta_*^{(s)}(\exp(z))$  along any horizontal line  $y = \pm 2k\pi + \varepsilon$ , k an integer, and the proof of Lemma 1 implies that there will exist a positive real number a such that  $z^r \eta_*^{(s)}(\exp(z))$  will have no zero on or to the right of the vertical line x = a. Consider the rectangle  $R_{ka}$  defined by  $0 \le x \le a$ ,  $-2k\pi + \varepsilon \le y \le 2k\pi + \varepsilon$ . Then the angular variation of  $\omega$ , as z tranverses the upper, lower and right sides of  $R_{ka}$ , will differ from that of  $\omega = z^r \eta_*^{(s)}(\exp(z))$  only by a number  $\delta_2$  where  $\delta_2 \rightarrow 0$  as  $k \rightarrow \infty$ . Since the boundary of  $R_{ka}$  has been shown to be free of poles and zeros, Cauchy's index theorem easily implies the desired result for  $\omega$ , and hence for  $\omega$ .

The following lemmas will complete the work preparatory to the final results.

Proof. From the definition of  $_{\omega}^{*}(a, b)$ , it is clear that  $_{\omega}^{*}(a, b)$ =  $_{\omega}^{*}(a, c) + _{\omega}^{*}(c, b)$  for any c. It is also clear from (3.23)
that for |a| large,  $_{\omega}^{*}(a, a + _{\varepsilon})$  differs little from the angular variation due to  $_{\omega}^{*}(a, a + _{\varepsilon})$  and  $_{\omega}^{*}(a, a + _{\varepsilon})$  and  $_{\omega}^{*}(a, a + _{\varepsilon})$  differs little from the angular variation due to  $_{\omega}^{*}(a, a + _{\varepsilon})$  and  $_{\omega}^{*}(a, a + _{\varepsilon})$  differs little from the angular variation due to  $_{\omega}^{*}(a, a + _{\varepsilon})$  and  $_{\omega}^{*}(a, a + _{\varepsilon})$  differs little from the angular variation due to  $_{\omega}^{*}(a, a + _{\varepsilon})$  and  $_{\omega}^{*}(a, a + _{\varepsilon})$  differs little from the angular variation  $_{\omega}^{*}(a, a + _{\varepsilon})$  and  $_{\omega}^{*}(a, a + _{\varepsilon})$  and  $_{\omega}^{*}(a, a + _{\varepsilon})$  differs little from the angular variation due to  $_{\omega}^{*}(a, a + _{\varepsilon})$  and  $_{\omega}^{*}(a, a + _{\varepsilon})$  differs little from the angular variation  $_{\omega}^{*}(a, a + _{\varepsilon})$  and  $_{\omega}^{*}(a, a + _{\varepsilon})$  differs little from the angular variation  $_{\omega}^{*}(a, a + _{\varepsilon})$  and  $_{\omega}^{*}(a, a + _{\varepsilon})$  differs little from the angular variation  $_{\omega}^{*}(a, a + _{\varepsilon})$  and  $_{\omega}^{*}(a, a + _{\varepsilon})$  differs  $_{\omega}^{*}(a, a + _{\varepsilon})$  and  $_{\omega}^{*}(a, a + _{\varepsilon})$  differs  $_{\omega}$ 

$$\omega^*(a, a + \epsilon) = \sum_{n=0}^{s} n\epsilon + \delta_4$$

where  $\delta_4 \to 0$  as  $a \to \pm \infty$ . Similarly,

$$\omega^*(b, b + \varepsilon) = \sum_{n=0}^{\infty} n\varepsilon + \delta_5$$

where  $\delta_5 \rightarrow 0$  as  $b \rightarrow \pm \infty$ . Then

$$\omega^*(a + \varepsilon, b + \varepsilon) = \omega^*(a, b) - \omega^*(a, a+\varepsilon) + \omega^*(b, b+\varepsilon)$$

which implies

$$w^*(a + \epsilon, b + \epsilon) = w^*(a, b) + \delta_3$$

where  $\delta_3 \to 0$  as  $a \to \pm \infty$  and  $b \to \pm \infty$ . Q.E.D.

- Lemma 3.4. Let h(z,t) have a principal term  $a_{rs}z^{r}t^{s}$ , and f(z,u,v) and g(z,u,v) have principal terms  $z^{r}(s)(u,v)$  and  $z^{r}(u,v)$  respectively. Then for  $\lambda$  and  $\mu$  arbitrary real numbers not both zero, there exists a number  $\varepsilon$  such that  $\lambda \varphi^{(s)}(\varepsilon + iy) + \mu \psi^{(s)}(\varepsilon + iy) \neq 0$  for all real y.
- Proof. By definition,  $\lambda \varphi_*^{-(s)}(z) + \psi_*^{-(s)}(z)$  is a polynomial in cos z and sin z and if not identically zero for all z, Lemma 2 will imply the desired result immediately. If either  $\varphi_*^{(s)}(z)$  or  $\psi_*^{-(s)}(z)$  were zero for all z, an appropriate choice of  $\lambda$  and  $\mu$  could be made such that the polynomial would be always zero. However, the proof of Theorem 3.2 clearly shows that neither  $\varphi_*^{(s)}(z)$  nor  $\varphi_*^{(s)}(z)$  can be zero for all z. The remaining possibility is for  $\varphi_*^{(s)}(z)$  and  $\varphi_*^{(s)}(z)$  to satisfy

$$-\frac{(s)}{\varphi_{*}}(z) = -\mu/\lambda - \frac{(s)}{\psi_{*}}(z)$$
 (3.25)

for some  $\lambda$  and  $\mu$ , neither zero, and all z. This could obviously occur if  $\phi_f^{(n)}(\cos z, \sin z)$  and  $\psi_r^{(n)}(\cos z, \sin z)$ 

were non-zero only for n=0, and this would imply a principal term for H(z) of the form  $a_{r0}^{\ \ z}$ . But this can occur only if H(z) is degenerate, a contradiction to a fundamental hypothesis. With s now shown to be non-zero, the proof of Theorem 3.2 is again examined, and it is noted that (3.18), (3.19) and (3.20) imply that for (3.25) to hold, it is necessary that

$$\varphi_{\rm r}^{(n)}(\cos z, \sin z) = -\mu/\lambda \psi_{\rm r}^{(n)}(\cos z, \sin z)$$
 (3.26)

for all  $n \le s$ . But this implies that

Re { 
$$a_{rn}^{i}(\cos(nz) + i \sin(sz))$$
 }  
=  $-\mu/\lambda \text{ Im } \{a_{rn}^{i}(\cos(nz) + i\sin(nz))\}$  (3.27)

for all  $n \le s$ . For any value of r, however, (3.27) can be shown to imply that  $a_{rs} = 0$ , a final contradiction. Q.E.D.

Lemma 3.5. Let h(z,t) have a principal term and write H(iy = F(y) + i G(y)). Suppose  $\Delta \omega (-2k\pi, 2k\pi) = 4sk\pi + \pi r + \delta_6$  where  $\Delta = \pm 1$  and  $-\pi/2 \le \delta_6 \le \pi/2$ . Then for  $\lambda$  and  $\mu$  arbitrary real numbers not both zero,  $\lambda F(y) + \mu G(y)$  has only real and simple zeros and

$$\Lambda [G'(y) F(y) - G(y) F'(y)] > 0$$
 (3.28)

for all y.

Proof. Let  $\lambda$  and  $\mu$  be given and consider the curve traced by the vector  $\omega = H(iy)$  as it subtends an angle within  $\pi/2$  radians of  $4sk_{\Pi}+_{\Pi}r$  radians in the positive or negative direction as y varies from  $-2k_{\Pi}$  to  $2k_{\Pi}$ . This curve intersects the line  $\lambda\omega' + \mu\omega'' = 0$  ( $\omega' + i\omega'' = \omega$ ) for at least 4ks+r distinct values y, which implies that  $\lambda F(y) + \mu G(y)$  has at least 4ks+r real zeros for  $-2k_{\Pi} \le y \le 2k_{\Pi}$ . By Lemma 3.3, the same result holds for  $-2k_{\Pi} + \varepsilon \le y \le 2k_{\Pi} + \varepsilon$ , any  $\varepsilon$ , if k is sufficiently large.

By Lemma 3.4, choose an  $\varepsilon$  such that  $\lambda \overset{-}{\phi} (\varepsilon + iy) + \overset{-}{\mu} (\varepsilon) (\varepsilon + iy)$  is non-zero for all real y. Then by Theorem 3.2,  $\lambda F(y) + \mu G(y)$  will have, for sufficiently large k, no more than 4sk+r zeros for  $-2k\pi+\varepsilon \le y \le 2k\pi+\varepsilon$ , and hence all the zeros are real and simple. The simplicity of the zeros implies that the curve traced by  $\omega$  always moves in a constant direction, and since the velocity of rotation is given by

$$\frac{d_w}{dy}^* = \frac{G'(y) F(y) - G(y) F'(y)}{F^2(y) + G^2(y)},$$

clearly

$$\Delta[G'(y) F(y) - G(y) F'(y)] > 0$$
 for all y. Q.E.D.

The final results are now presented.

Theorem 3.5. Suppose all the zeros of H(z) are strictly in the left half plane, and write H(iy) = F(y) + iG(y). Then the zeros of F(y) and G(y) are real, simple and alternate and G'(y) F(y) - F'(y) G(y) > 0 for all y.

Proof. The contrapositive of Theorem 3.1 implies that h(z,t) has a principal term. By Lemma 3.1, choose an  $\varepsilon$  such that  $\eta_*^{(s)}(\exp(x+i~(2n\pi+\varepsilon)))$  is non-zero for all positive real x and integer n. Applying Theorem 3.4, and since  $N_k=0$ ,

$$\omega^* \left(-2k_{\pi} + \epsilon, 2k_{\pi} + \epsilon\right) = 4k_{\pi} + \pi r + \delta'_{k}$$

where  $\delta_k' \to 0$  as  $k \to \infty$ . For  $\chi = 1$  and  $\mu = 0$ , Lemmas 3.3 and 3.5 imply that the zeros of F(y) are real and simple. For  $\chi = 0$  and  $\mu = 1$ , the zeros of G(y) are similarly real and simple. Lemma 3.5 also implies that

$$F(y) G'(y) - F'(y) G(y) > 0$$

for all real y, which, in turn, implies that the zeros of F(y) and G(y) alternate. Q.E.D.

- Theorem 3.6. Let H(z) be non-degenerate with no pole at the origin and write H(iy) = r(y) + i G(y). Then in order for all of the zeros of H(z) to lie strictly in the left half plane, each of the following conditions is both necessary and sufficient:
  - (i) All of the zeros of F(y) and G(y) are real and alternate and F(y) G'(y) F'(y) G(y) > 0 for some y.

    (ii) All of the zeros of F(y) are real and at each zero,  $Y_0$ , of F(y),  $F'(y_0)$   $G(y_0) < 0$ .

    (iii) All the zeros of G(y) are real and at each zero,  $Y_0$ , of G(y),  $F(y_0)$   $G'(y_0) > 0$ .

Proof. Necessity.

The necessity of each condition is immediately obvious from a consideration of Theorem 3.5.

Sufficiency.

(i). Since the zeros of F(y) and G(y) are real, the contrapositive of Theorem 3.3 implies that f(z,u,v) and g(z,u,v) both have principal terms, and hence so does h(z,t). By Lemma 3.2, an  $\varepsilon$  exists completing the hypothesis of Theorem 3.2 and F(y) and G(y) thus each have 4sk+r real zeros for  $-2k\pi+\varepsilon \le y \le 2\pi+\varepsilon$ , k sufficiently large. The rotation of H(iy) is thus at least  $\pm (rks\pi+\pi r) \pm \pi/2$  radians and hence

$$\Delta \omega^* (-2k\pi + \epsilon, 2k\pi + \epsilon) \ge 4ks\pi + \pi r - \pi/2.$$

Then by Lemma 3.3,

$$\Delta \omega^* (-2k\pi, 2k\pi) \ge 4ks\pi + \pi r - \pi/2$$

for k sufficiently large, and the results of Lemma 3.5 obviously hold implying

$$\Delta[F(y) G'(y) - F'(y) G(y)] > 0$$

for all y. Since F(y) G'(y) - F'(y) G(y) > 0 for some y by hypothesis,  $\Delta = +1$ . Thus,

$$\omega^*(-2k\pi, 2k\pi) \ge 4sk\pi + \pi r - \pi/2$$
 (3.29)

Now since the zeros of F(y) and G(y) alternate, they are simple and H(z) has no imaginary zero. Use of Lemma 3.1 completes the hypothesis of Theorem 3.4, and

where 
$$\delta_{\mathbf{k}}^{*} \rightarrow 0$$
 as  $\mathbf{k} \rightarrow \infty$ . By Lemma 3.3,  
 $\mathbf{w}^{*}(-2\mathbf{k}\pi + \varepsilon, 2\mathbf{k}\pi + \varepsilon) = 2\pi(2\mathbf{s}\mathbf{k} - \mathbf{N}_{\mathbf{k}} + \mathbf{r}/2) + \delta_{\mathbf{k}}$   
where  $\delta_{\mathbf{k}}^{*} \rightarrow 0$  as  $\mathbf{k} \rightarrow \infty$ . By Lemma 3.3,  
 $\mathbf{w}^{*}(-2\mathbf{k}\pi, 2\mathbf{k}\pi) = 4\mathbf{s}\mathbf{k}\pi + \pi\mathbf{r} - 2\pi \mathbf{N}_{\mathbf{k}} + \delta_{\mathbf{k}}^{\dagger}$  (3.30)

where  $\delta_k^{\prime} \to 0$  as  $k \to \infty$ . Then for large k, (3.29) and (3.30) imply that  $N_k = 0$ . Q.E.D.

(ii) and (iii). Proof of the sufficiency of conditions (ii) and (iii) follows that of condition (i) closely, with the two inequalities insuring the absence of imaginary zeros and setting the direction of rotation of the vector H(iy).

Q.E.D.

The results of Theorem 3.6 have left only two problems to be solved before definitive statements can be made concerning the asymptotic stability of any particular system. The first of these, illustrated in some detail in a following chapter, is the analytical difficulty involved in parametrically satisfying one of the three conditions. The second, mentioned earlier in this work, concerns the insufficiency of left half plane zero locations in implying global asymptotic stability. The following lemma will answer certain questions concerning this problem.

Lemma 3.6. Let  $P(z) = \mathcal{L} \{ p(t) \}$  and  $p(t) = \mathcal{L}^{-1} \{ P(z) \}$  be a well-defined Laplace transform pair and suppose all poles of P(z) lie to the left of the line  $R_{e}\{z\} = -c$ , where c is a strictly positive real number. Then

$$\lim_{t\to\infty} p(t) = 0.$$

Proof. By definition, p (t) is given by

$$p(t) = \frac{1}{2\pi j} \quad \underset{y \to \infty}{\text{Lim}} \quad \int_{z}^{\sigma + jy} P(z) e^{zt} dz$$

where the path of integration is any line  $z=\sigma$ , a constant damping insuring the convergence of P(z). Clearly  $\sigma=-c/2$  will suffice, and thus

$$p(t) = \frac{1}{2\pi j} \lim_{y \to \infty} \int_{-\frac{C}{2} - jy} P(z)e^{zt}dz.$$

Since the integration is for  $z = -c/2 + j_{\omega}$ ,  $-\infty < \infty < \infty$ , write

$$p(t) = \frac{1}{2\pi j} \lim_{y \to \infty} \int P(-\frac{c}{2} + j_{\omega}) = \frac{-ct}{2} e^{j\omega t} d(j_{\omega})$$
$$-\frac{c}{2} - jy$$

$$-\frac{c}{2}+jy$$
 and 
$$|p(t)| \leq \frac{1}{2\pi} \lim_{y \to \infty} \int |P(-\frac{c}{2}+j_{w})| e^{-\frac{ct}{2}} d_{w}.$$
 
$$-\frac{c}{2}-jy$$

But this implies

$$\begin{array}{c|c} \left| \; p(t) \; \right| \; \leq \; e^{\displaystyle \frac{-\frac{\mathbf{C}}{2}}{2} \; t} \; \underset{y \; \rightarrow \; \infty}{\underline{1}} \quad \begin{array}{c} -\frac{\mathbf{C}}{2} \; + j y \\ \int \left| \; P(-\frac{\mathbf{C}}{2} \; + \; j_{\text{\tiny $W$}}) \; \; \right| \; d_{\text{\tiny $W$}}, \\ -\frac{\mathbf{C}}{2} \; - j y \end{array}$$

and since the path of integration is free of singularities of P(z), the result immediately follows. Q.E.D.

This lemma implies the following theorem.


Theorem 3.7. Given the system of (1.6) - (1.7), suppose the determinantal equation is expressable as (3.6) with the usual aforementioned restrictions. Then the solutions of (1.6)-(1.7) will approach zero as  $t \to \infty$  if there exists a real number e > 0 such that H(z - e) has all its zeros strictly in the left half plane.

# IV. CHARACTERISTICS OF MATHEMATICAL MODEL SOLUTIONS Qualitative Characteristics

- E. C. Bratt ([17], p. 204) in summarizing the contributions of the Austrian school, states that the "...monetary overinvestment theorists have shown little interest in statistical verification." Rothbard ([10], pp. 3-5), a major Austrian supporter, believes that "economic theories cannot be 'tested' by historical or statistical fact.... Theory cannot emerge, Phoenix-like, from a cauldron of statistics; neither can statistics be used to test an economic theory." This dearth of factual numerical support for the Austrian theory makes difficult the testing of the system mathematical model, yet it can be shown that the model does admit to solutions which agree qualitatively with the Austrian conclusions. These conclusions may be summarized as follows:
  - (i) The boom period could continue indefinitely if credit could be expanded indefinitely ([8], p. 332).
  - (ii) The capital-goods industries are capable of fluctuating more widely than the consumer-goods industries ([10], pp. 16-17).
  - (iii) The capital-goods industries expand, in general, at the expense of expansion in the consumer-goods sector, leading to an out-of-phase cyclical behavior ([8], pp. 332-3).

Figure 2 illustrates a particular numerical example whose solution agrees with conclusion (i). Although this particular conclusion is often attacked as "...seriously in error" ([8], pp. 332-3), it is a basic tenet of the Austrian theory and any valid model must admit to this type solution. Solutions satisfying conclusion (ii) are easily obtained, merely increasing the damping effects on v(t) and sharply truncating the credit expansion input, as illustrated in Figure 3. Conclusion (iii) also is often the subject of criticism, since at first glance it seems to deny the possibility of depressed conditions existing in both industrial sectors. Figure 4, however, illustrates the proper implications of this conclusion, since the


## THE CONTINUING BOOM



$$x(t) = g(t) - \int_{0}^{1} 0.5 x(t-\tau) d\tau - 1.5 v(t)$$
 \_\_\_\_\_ g(t)  
$$v'(t) = -2.0 v(t) - 1.5 x(t), v(0) = 0$$
 \_\_\_\_\_ v(t)

Figure 2

#### THE CAPITAL-GOODS FLUCTUATION




$$x(t) = g(t) - \int_{0}^{1} 0.5 x(t - \tau) d\tau - 1.0 v(t)$$
 g(t)

$$v'(t) = -3.0 v(t) - 1.0 x(t), v(0) = 0$$
 \_\_\_\_ v(t)

Figure 3

## THE CYCLICAL SOLUTION



$$x(t) = g(t) - \int_0^t 0.5 \times (t - \tau) d\tau - 0.3 v(t)$$
 \_\_\_\_\_ g(t)

$$v'(t) = -2.0 v(t) - 0.5 x(t), v(0) = 0$$
 \_\_\_\_ v(t)

Figure 4

capital-goods investment might become rapidly depressed before the consumer-goods investment could recover. This implies an overall depressed period followed by the out-of-phase cyclical behavior as the system returns to its normal (zero) solution.


One final solution type concerns the existence of non-zero constant trajectories under zero input conditions. Prior to the Great Depression, the economy was considered a dynamic, always changing phenomenon. A constantly depressed economy, for example, was considered an impossibility. The events of the thirties proved an embarrassing counterexample, as year after year idle men loafed next to idle machines. It was John Maynard Keynes [18] who explained this paradox, simply noting that a depressed economy could be in perfect economic balance even though in the depths of social agony. Thus, any business cycle theory must admit solutions which exhibit this economic balance/social imbalance characteristic. Figure 5 illustrates such a constant solution, with a continuing investment in the capital-goods industries above the norm and a corresponding continuing investment in consumer-goods industries below the norm. This, of course, corresponds to a continually inflated economy with no continuing credit expansion.

## Quantitative Characteristics

As previously mentioned, the quantitative aspects of the Austrian theory are not presented by its proponents. While the required statistics are available for individual industries, the Austrian theorists have not attempted the dichotomization and compilation of such statistics necessary to support their conjectures. The following is an attempt to characterize the cyclical patterns of the two investment sectors (capital-goods and consumer goods) by two specific quantities, the value of building permits and department store sales, in a brief examination of the minor business cycle of 1921-1924.

The value of building permits issued was used to characterize the investment in capital-goods industries. Since the construction industry

## A CONSTANT SOLUTION



$$x(t) = g(t) - \int_{0}^{1} 0.5 x(t - \tau) d\tau - 2.0 v(t) \qquad g(t)$$

$$v'(t) = -2.0 v(t) - 1.5 x(t), v(0) = 0$$

$$v(t)$$

Figure 5

exhibits long delays and is historically hurt by tight-money situations, it should approximate to some degree the overall investment nature of the capital-goods industries. To approximate the consumer-goods sector, retail department store sales was chosen. As the department stores vary their stock levels to compensate for sales fluctuations, a rapid, short term investment process is carried out, typical of investment in consumer-goods industries.

The raw data for the value of building permits issued and department store sales (from [19], pp. 194-6) is presented in Figures 6 and 7, with the trend chosen in each case as a linear least squares fit. The governmental credit policies are presented in Figure 8, compiled from Rothbard([10], pp. 101-5). In the mathematical model, the influence function was chosen as a constant for eighteen months duration, and the influence function amplitude  $\alpha$ ,  $\beta$ , and  $\gamma$  varied until the model solution minimized a least squares error criterion for the period January, 1921, to July, 1922. With the parameters then fixed at  $\rho = 0.5$ ,  $\alpha = 0.5$ ,  $\beta = 3.5$  and  $\gamma = 1.0$ , the solution was extended under the actual credit policy input to January, 1924. The model solution is compared with the actual values in Figures 9 and 10.

The mathematical model solution does not fit the actual quantities well in two respects; the building slump from January to July, 1922, is not fitted and the rising department store sales after July, 1922, is lagged by several months. It should be noted, however, that the Austrian theory itself does not explain these results. Since government credit expansion monotonically increased until July, 1922, neither a drop in building nor an increase in consumer demand could be predicted. Either the Austrian theory is at fault or, more likely, the use of relatively small portions of the economic process to characterize larger segments is of limited accuracy. For example, factors such as lumber prices and weather could cause fluctuations in the construction industry but would not greatly affect the overall capital-goods investment

# INDEX OF VALUE OF BUILDING PERMITS JANUARY 1921 - JANUARY 1924

(July 1921 = 100)




Figure 6

# DEPARTMENT STORE SALES JANUARY 1921 - JANUARY 1924

(July 1921 = 100)




Figure 7

# GOVERNMENT CREDIT EXPANSION IN MILLIONS OF DOLLARS JANUARY 1921 - JANUARY 1924




Figure 8

# VALUE OF BUILDING PERMITS DEVIATION FROM THE TREND: IN PERCENT OF JULY 1921 VALUE




Figure 9

# DEPARTMENT STORE SALES DEVIATION FROM THE TREND: IN PERCENT OF JULY 1921 SALES




Figure 10

process. But even with these inaccuracies, a prognostic solution under proposed credit policies would, in July of 1922, have indicated the results of such a precipitous drop in monetary supply.

#### V. A ZERO LOCATION PROBLEM

With some correspondence having been established between the theory conclusions and the model solutions, this chapter will present a specific numerical example illustrating the numerical applications of Theorem 3.6. Suppose the system equations are given by

$$x(t) = g(t) - \int_{0}^{1} 0.5 x (t-\tau) d\tau - \alpha v(t)$$

$$v'(t) = -2.0 v(t) - \gamma x(t), v(0) = 0,$$

and the system stability is to be studied as a function of the coupling parameters  $\alpha$  and  $\gamma$ . The zero locations are determined by

$$H(z) = [z + (2.5 - \alpha \gamma) + 1/z] e^{z} - 0.5 - 1/z,$$

and condition (i) will be utilized. Let z = x + iy,  $\xi = 2.5 - \alpha \gamma$  and

$$H(iy) = [i(y - 1/y) + \xi][\cos y + i\sin y] - 0.5 + 1/y,$$

which implies

$$F(y) = \xi \cos y - (y - 1/y) \sin y - 0.5$$

$$G(y) = \xi \sin y + (y - 1/y) \cos y + 1/y$$
.

The zeros of F(y) will be examined first, utilizing the fundamental Theorem 3.2. The principal term of F(y) is-ysiny, hence

$$\varphi$$
 (s) (cosy, siny) = - siny

and

$$\varphi_{\star}^{-(s)}(z) = -\sin z.$$

Then

$$\varphi_{\star}^{-(s)}(\varepsilon + 2n\pi + iy) = -\sin(\varepsilon + 2n\pi + iy)$$

is clearly non-zero for all real y and integer n if  $\varepsilon = \pi/2$ . It must now be shown that F(y) has, for sufficiently large k, exactly 4k + 1 zeros (s = 1 in this case) for  $-2k\pi + \pi/2 \le y \le 2k\pi + \pi/2$ . Figure 11 graphically illustrates the following observations:

## THE ZEROS OF F(y)




Figure 11

- (i) For  $\xi$  less than -0.5, F(y) will have no zero for  $0 \le y \le \pi/2$ .
- (ii) For  $\xi$  exactly -0.5, F(y) will have a double zero at y = 0.
- (iii) For  $\xi$  greater than -0.5, F(y) will have exactly one zero for  $0 < y \le \pi/2$ .
- (iv) For  $\xi \ge -0.5$ , F(y) will have exactly one zero in each open interval  $n\pi + \pi/2 < y < (n+1)\pi + \pi/2$ ,  $n=1,2,\ldots$

The preceding observations, and noting that (y-1/y) siny and cosy -0.5 are even functions of y, imply that F(y) has, for sufficiently large k, exactly 4k+1 zeros for  $-2k\pi+\pi/2 \le y \le 2k\pi+\pi/2$  if and only if  $\xi \ge -0.5$ . The zeros of G(y) are next examined.

The principal term of G(y) is ycosy, and thus

$$\psi_{+}^{-(s)}(\varepsilon + 2n\pi + iy) = \cos(\varepsilon + 2n\pi + iy)$$

is non-zero for all y and integer n if  $\varepsilon = 0$ . Noting that  $\xi \ge -0.5$  has already been shown to be a necessary condition, it may now be examined only as a sufficiency requirement. Figure 12 illustrates the following:

- (i) For  $\xi \ge -0.5$ , G(y) has exactly one zero in each interval  $n\pi < y \le (n+1)\pi$ , n = 0,1,2,...
- (ii) For any  $\xi$ , G(y) has a zero at y=0, and that zero is simple. The preceding observations, and noting that  $\xi \sin y$  and  $-1/y (y 1/y)\cos y$  are both odd functions of y, imply that  $\xi \ge -0.5$  is sufficient to insure that G(y) has, for sufficiently large k, exactly 4k + 1 zeros for  $-2k\pi \le y \le 2k\pi$ .

The relative positions of the zeros of F(y) and G(y) must next be examined. It is first noted that for  $\xi = -0.5$ , F(y) has a double root at the origin. Since this obviously would not allow the zeros of F(y) and G(y) to alternate, the condition must be changed to  $\xi > -0.5$ . Suppose now that  $\xi > 0.5$ . The zero locations of F(y) may then be described by




Figure 12

$$0 < y_{10} < \pi/2$$

$$y_{20} = \pi + \theta_{2}$$

$$\vdots$$

$$(n-1)\pi \le y_{n0} < (n-1)\pi + \pi/2, n \text{ odd}$$

$$y_{n0} = (n-1)\pi + \theta_{n}, n \text{ even}$$

$$\vdots$$

where  $0 < \theta_n < \pi/2$  and  $\theta_n$  is defined by

$$\xi \cos((n-1)\pi + \theta_n) - 0.5 = ((n-1)\pi + \theta_n) - 1/((n-1)\pi + \theta_n) \sin((n+1)\pi + \theta_n)$$
, n even.

Similarly, the zero locations for G(y) are given by

$$y'_{10} = 0$$
 $\pi/2 < y'_{20} < \pi$ 
 $y'_{30} = \pi + \theta_{3}'$ 
 $\vdots$ 
 $(n-2)\pi + \pi/2 < y'_{n0} < (n-1)\pi$ , n even

 $y'_{n0} = (n-2)\pi + \theta_{n}'$ , n odd,

where  $\theta$  is defined by

$$\xi \sin((n-2)\pi + \theta_n') = -1/((n-2)\pi + \theta_n') - ((n-2)\pi + \theta_n') - ((n-2)\pi + \theta_n') - 1/((n-2)\pi + \theta_n'))\cos((n-2)\pi + \theta_n'), \text{ n odd.}$$

It is clear that if  $\theta_n'>\pi/2$  ,  $y_{n0}'$  lies to the right of  $y_{n-1}$  ,0. If  $\theta_n'<\pi/2$  , it must be shown that

$$\theta_n < \theta_n$$

for all appropriate n. While a lengthy and complicated analysis is required to prove this result rigorously, the following geometric argument can be briefly presented. A short study of Figure 11 shows that  $\theta_n$  will be large (slightly less than  $\pi/2$  radians) only if  $\xi$  is in some sense large and y in

some sense small. Figure 12, however, indicates that these same conditions result in  $\theta_n$  being very large, approaching  $\pi$  radians. On the other hand, a brief calculation shows that  $\theta_n$  is at least 1.5 radians, and the minimal values occur for  $\xi$  small. But if  $\xi$  is small, the zeros of F(y) lie only slightly to the right of odd multiples of  $\pi$ , implying that  $\theta_n$  is indeed less than 1.5 radians.

Similar (but somewhat simpler) arguments for  $-0.5 < \xi \le 0.5$  show that  $\xi > -0.5$  is indeed a sufficient condition for the zeros of F(y) and G(y) to alternate. For the final step, it must be shown that  $\xi > -0.5$  implies that

$$F(y)G'(y) - F'(y)G(y) > 0$$

for some y. The appropriate derivatives are given by

$$F'(y) = -a \sin y - (y - 1/y) \cos y - (1 + 1/y^{2}) \sin y,$$

$$G'(y) = a \cos y - (y - 1/y) \sin y + (1 + 1/y^{2}) \cos y - 1/y^{2}.$$

Suppose y = 0 is chosen. Then G(y) = 0,

$$F(0) = \xi + \lim_{y \to 0} \frac{\sin y}{y} - 0.5$$
  
=  $\xi + 0.5$ 

and

$$G'(0) = \xi + \lim_{y \to 0} \frac{\sin y}{y} + \lim_{y \to 0} \left\{ \frac{\cos y}{y^2} - \frac{1}{y^2} \right\} + 1$$
$$= \xi + 1.5.$$

Thus

$$F(0)G'(0) = (\xi + 0.5)(\xi + 1.5)$$

and F(0)G'(0) > 0 if  $\xi > -0.5$ . Finally, then, the zeros are all strictly in the left half plane if and only if  $\alpha \gamma < 3.0$ .

#### VI. CONCLUSIONS

Evans ([8], p. 415), in speaking of the multiplier-accelerator econometric model, states that "...second-order difference equations are no longer adequate to determine the solution in models with more realistic and complicated lag structures." To overcome these problems, he suggests that an appropriate approach would be to "...state a more general theory of the cycle, estimate this theory empirically, and examine its behavior by actual simulation." While it might certainly be true that a more general theory of economy is needed, it is not so clear that any such new theory should be examined from an empirical point of view. It is certainly possible that a mathematical structure more complex than second-order difference equations might very accurately model such a new theory.

To support this conjecture, this work has presented a new modelling form, the coupled renewal-differential equation system, and with this structure modelled the Austrian theory of business cycles.

Mathematical results presented include conditions insuring the existence, uniqueness and Laplace transformability of solutions, and an illustration of the utilization of a differential equation solving subroutine in calculating actual numerical solutions. Stability results include a generalization of results due to Pontryagin [13] which gives necessary and sufficient conditions for all the system poles to lie strictly in the left half plane. Further numerical results include a qualitative comparison of the Austrian theory conclusions and the types of solutions the mathematical model possesses, and finally an illustration of the numerical use of the stability conditions.

While the Austrian theory of business cycles has been the only economic theory modelled, no attempt has been made to either support or reject, through the numerical results, the conclusions of the Austrian school. This particular theory was used only in an illustrative capacity to indicate the possible use of the modelling structure.

#### BIBLIOGRAPHY

- [1] Lotka, A. J., "A Contribution to the Theory of Self-renewing Aggregates with Special Reference to Industrial Replacement," Ann. Math. Statistics, Vol. 10, 1939.
- [2] Wangersky, P. J. and Cunningham, W. J., "On Time Lags in Equations of Growth," Proc. Nat. Acad. of Sci., Vol. 46, 1956.
- [3] Lotka, A. J., <u>Theorie Analytique des Associationes Biologiques</u>, Hermann et Cie, Paris, 1939.
- [4] Fein, A. E., "Influence of a Variable Ejection Probability on the Displacement of Atoms," Phy. Rev., Vol. 109, 1958.
- [5] Weibel, E. S., "Dynamic Stabilization of a Plasma Column," Space Technology Laboratory, 1960.
- [6] Melzak, Z. A., "Some Mathematical Problems in Retrograde Nerve Degeneration," J. Math. Anal. Appl., Vol. 2, 1961.
- [7] Heilbroner, R. L., <u>The Worldly Philosophers</u>, Simon and Schuster, New York, 1961.
- [8] Evans, M. K., <u>Macroeconomic Theory</u>, Harper and Row, New York, 1969.
- [9] Smith, Adam, Wealth of Nations, Henry Regnery, Chicago, 1953.
- [10] Rothbard, M. N., <u>America's Great Depression</u>, D. Van Nostrand, Princeton, 1963.
- [11] Mises, L. von, <u>Human Action</u>, Yale University Press, New Haven, 1949.
- [12] Hayek, F. A., <u>Prices and Production</u>, 2nd ed., George Routledge, London, 1935.
- [13] Pontryagin, L. S., "On the Zeros of Some Elementary Transcendental Functions," Am. Math. Soc. Trans., Series 2, Vol. 1, 1955.
- [14] Bellman, R. and Cooke, K. L., <u>Differential-Difference Equations</u>, Academic Press, New York, 1963.

- [15] Nordsieck, A., <u>Mathematics of Computation</u>, Vol. XVI, No. 77, 1962.
- [16] Khinchin, A. I., <u>A Course of Mathematical Analysis</u>, Hindustan, India, 1960 (Transl. from Russian).
- [17] Bratt, E. C., <u>Business Cycles and Forecasting</u>, Richard D. Irwin, Homewood, Ill., 1953.
- [18] Keynes, J. M., The General Theory of Employment, Interest and Money, Harcourt, Brace and Co., New York, 1936.
- [19] Lee, M. W., Economic Fluctuations, Richard D. Irwin, Homewood, Ill., 1955.

#### DISTRIBUTION LIST\*

#### Grant No. AF-AFOSR 766-67

Mod. No. AFOSR-67-0766E

Report on the Joint Services Electronics Program for period ending 30 September 1969

#### DEPARTMENT OF DEFENSE

Dr. A. A. Dougal (1) Asst Director (Research) Ofc of Defense Res & Eng Department of Defense Washington, D. C. 20301

Office of Deputy Director (1) (Research and Information, Rm 3D1037 Department of Defense The Pentagon Washington, D. C. 20301

Director (1) Advanced Research Projects Agency Department of Defense Washington, D. C. 20301

Director for Materials Sciences (1) Advanced Research Projects Agency Department of Defense Washington, D. C. 20301

Headquarters (1) Defense Communications Agency (340) Washington, D. C. 20305

Defense Documentation Center (50) Attn: DDC-TCA Cameron Station Alexandria, Virginia 22314

Director (1) National Security Agency Attn: TDL Fort George G. Meade, Md. 20755

Weapons Systems Evaluation Group (1) Attn: Colonel Blaine O. Vogt 400 Army-Navy Drive Arlington, Virginia 22202

Central Intelligence Agency (1) Attn: OCR/DD Publications Washington, D. C. 20505

#### DEPARTMENT OF THE AIR FORCE

Hq USAF (AFRDDD) (1) The Pentagon Washington, D. C. 20330

Hq USAF (AFRDDG) (1) The Pentagon Washington, D. C. 20330

Hq USAF (AFRDSD) (1) The Pentagon Washington, D. C. 20330

Colonel E. P. Gaines, Jr. (1) ACDA/FO 1901 Pennsylvania Avenue N.W. Washington, D. C. 20451

Lt. Col. R. B. Kalisch (SREE) (5) Chief, Electronics Division Directorate of Engineering Sciences Air Force Office of Scientific Research Arlington, Virginia 22209

Dr. I. R. Mirman (1) AFSC (SCT) Andrews Air Force Base, Md. 20331

AFSC (SCTSE)(1) ws Air Force Base, Md. 20331

Mr. Morton M. Pavane, Chief (1) AFSC Scientific and Technical Listson Office 26 Federal Plaza, Suite 1313 New York, New York 10007

;

Rome Air Development Center (1) Attn: Documents Library (EMTLD)
Griffiss Air Force Base, New York 13440 Mr. H. E. Webb (EMIIS) (1) Rome Air Development Center Griffiss Air Force Base, New York 13440

Dr. L. M. Hollingsworth (1) AFCRL (CRN) L. G. Hanscom Field Bedford, Massachusetts 01730

AFCRL (CRMPLR), Stop 29 (1) AFCRL Research Library L. G. Hanscom Field Bedford . Massachusetts 01730

Hg ESD (ESTI) (2) L. G. Hanscom Field Bedford, Massachusetts 01730

Professor J. J. D'Azzo (1) Dept of Electrical Engineering Air Force Institute of Technology Wright-Patterson AFB, Ohio 45433

Dr. H. V. Noble (CAVT) (1) Air Force Avionics Laboratory Wright-Patterson AFB, Ohio 45433

Air Force Avionics Laboratory Wright-Patterson AFB, Ohio 45433

AFAL (AVTA/R. D. Larson) (1) Wright-Patterson AFB, Ohio 45433

Director of Faculty Research (1) Department of the Air Force U. S. Air Force Academy Colorado Springs, Colorado 80840

Academy Library (DFSLB) (1) USAF Academy Colorado Springs, Colorado 80840

Director (1) Aerospace Mechanics Division Frank J. Seiler Research Laboratory (OAR) USAF Academy Colorado Springs, Colorado 80840

Director, USAF PROJECT RAND (1) Via: Air Force Liaison Office the RAND Corporation Attn: Library D 1700 Main Street Santa Monica, California 90406

Hq SAMSO (SMTTA/Lt. Nelson) (1) AF Unit Post Office Los Angeles, California 90045

Det 6, Hq OAR (1) Air Force Unit Post Office Los Angeles, California 90045

AUL3T-9663 (1) Maxwell AFB, Alabama 36112

AFETR Technical Library (1) (ETV, MU-135) Patrick AFB, Florida 32925

ADTC (ADBPS-12) (1) Eglin AFB, Florida 32542

Mr. B. R. Locke (1) Technical Adviser, Requirements
USAF Security Service
Kelly Air Force Base, Texas 78241

Hg AMD (AMR) (1) ooks AFB, Texas 78235 USAFSAM (SMKOR) (1) Brooks AFB, Texas 78235

Commanding General (2) Attn: STEWS-RE-L, Technical Library White Sands Missile Range New Mexico 88002

Hq AEDC (AETS) (1) Attn: Library/Documents Arnold AFS, Tennessee 37389

European Office of Aerospace Research (1)
APO New York 09667

Electromagnetic Compatibility Analysis Center (1) (ECAC), Attn: ACLP North Severn Annapolis, Maryland 21402

#### DEPARTMENT OF THE ARMY

Physical & Engineering Sciences Division (1) U. S. Army Research Office 3045 Columbia Pike Arlington, Virginia 22204

Commanding General (1) U. S. Army Security Agency Attn: IARD-T Arlington Hall Station Arlington, Virginia 22212

Commanding General (1) U. S. Army Materiel Command Attn: AMCRD-TP Washington, D. C. 20315

Commanding Officer (1) Harry Diamond Laboratories
Attn: Dr. Berthold Altman (AMXDO-TI) Connecticut Avenue and Van Ness Street N.W Washington, D. C. 20438

Director (1) Walter Reed Army Institute of Research Walter Reed Army Medical Center Washington, D. C. 20012

Commanding Officer (AMXRD-BAT) (1) U. S. Army Ballistics Research Laboratory Aberdeen Proving Ground Aberdeen, Maryland 21005

Technical Director (1) U. S. Army Limited War Laboratory Aberdeen Proving Ground Aberdeen, Maryland 21005

Commanding Officer (1) Human Engineering Laboratories Aberdeen Proving Ground Aberdeen, Maryland 21005

U. S. Army Munitions Command (1) Attn: Science & Technology Information Branch, Bldg. 59 Piontinny Arsenal, SMUPA-VA6 Dover, New Jersey 07801

U. S. Army Mobility Equipment Research and Development Center (1) Attn: Technical Document Center Bldg. 315

Fort Belvoir, Virginia 22060

Director (1) U. S. Army Engineer Geodesy, Intelligence & Mapping Research and Development Agency Fort Belvoir, Virginia 22060

Dr. Herman Robl (1) Deputy Chief Scientist U. S. Army Research Office (Durham) Box CM, Duke Station Durham, North Carolina 27706

Richard O. Ulsh (CRDARD-IPO) (1) U. S. Army Research Office (Durham) Box CM, Duke Station Durham, North Carolina 27706

Technical Director (SMUFA-A2000-107-1) (1) Frankford Arsenal Philadelphia, Pennsylvania 19137

Redstone Scientific Information Attn: Chief, Document Section U.S. Army Missile Command Redstone Arsenal, Alabama 35809

Commanding General (1) U. S. Army Missile Command Attn: AMSMI-REX Redstone Arsensi, Alabama 35809

Commanding General (1) U. S. Army Strategic Communications Command Attn: SCC-CG-SAE Fort Huachuca, Arizona 85613

Commanding Officer (1) Army Materials and Mechanics Research Center Attn: Dr. H. Priest Watertown Arsenal Watertown, Massachusetts 02172

Commandant (1) U. S. Army Air Defense School Attn: Missile Science Division, C & S Dept P.O. Box 9390 Fort Bliss, Texas 79916

Commandant (1) U. S. Army Command and General Staff College Attn: Acquisitions, Lib Div Fort Leavenworth, Kansas 66027

Commanding Officer (1) U.S. Army Electronics R&D Activity White Sands Missile Range New Mexico 88002

Mr. Norman J. Field, AMSEL-RD-S (1) Chief, Office of Science & Technology Research and Development Directorate U. S. Army Electronics Command Fort Monmouth, New Jersey 07703

Mr. Robert O. Parker, AMSEL-RD-S (1) Executive Secretary, JSTAC U. S. Army Electronics Command Fort Monmouth, New Jersey 07703

Commanding General U.S. Army Electronics Command Tor Monouth, New Jersey 07703

Attn: AMSEL-SC, RD-QF, RD-MT,

XL-D, XL-T, XL-C, XL-S(Dr. R. Buser),

HL-CT-DD, HL-CT-R, HL-CT-L(Dr. W.

S. McAfee), HL-CT-O, HL-CT-I, S. McAree), NI-CI-C, NI-CI-I,
HI-CT-A, NL-D, NI-A,
NL-P, NL-P-2 (Mr. D. Haratz)
NL-R (Mr. R. Kulinyi), NL-S, KL-D,
KL-E, KL-S(Dr. H. Jacoba), KL-SM
(Dra. Schiel/Hieslmatr), KL-T, VL-D, VL-F(Mr. R. J. Niemela), WL-D

\*The Joint Services Technical Advisory Committee has established this list for the regular distribution of reports on the electronics research program of The University of Texas at Austin. Additional addresses may be included on their written request to:

> Mr. Robert O. Parker (AMSEL-RD-S) U. S. Army Electronics Command Fort Monmouth, New Jersey 07703

An appropriate endorsement by a Department of Defense sponsor is required except on requests from a Federal Agency.

Dr. A. D. Schnitzler, AMSEL-HL-NVII (1) Night Vision Laboratory, USAECOM Fort Belvoir, Virginia 22060

Dr. G. M. Janney, AMSEL-HL-NVOR (1) Night Vision Laboratory, USAECOM Fort Belvoir, Virginia 22060

Atmospheric Sciences Office (1) Atmospheric Sciences Laboratory White Sands Missile Range New Mexico 88002

Missile Electronic Warfare
Technical Area, AMSEL-WT-MT (1)
White Sands Missile Range
New Mexico 88002

Deputy for Research and Engineering (AMSWE-DRE) (1) U. S. Army Weapons Command Rock Island Arsenal Rock Island, Illiniois 61201

Project Manager (1)
Common Positioning & Navigation
Systems
Attn: Harold H. Bahr

(AMCPM-NS-TM), Bldg. 439 U. S. Army Electronics Command Fort Monmouth, New Jersey 07703

Director (1)
U. S. Army Advanced Materiel
Concepts Agency
Washington, D. C. 20315

#### DEPARTMENT OF THE NAVY

Director, Electronic Programs (3) Attn: Code 427 Department of the Navy Washington, D. C. 20360

Commander (1) U.S. Naval Security Group Command Attn: G43 3801 Nebraska Avenue Washington, D. C. 20390

Director
Naval Research Laboratory
Washington, D. C. 20390
Attn: Code 2027 (6)
Dr. W.C. Hall, Code 7000 (1)
Dr. A. Brodzinsky,
Supt. Elec. Div. (1)

Dr. G. M. R. Winkler (1)
Director, Time Service Division
U. S. Naval Observatory
Washington, D. C. 20390

Naval Air Systems Command (2) AIR 03 Washington, D. C. 20360

Naval Ship Systems Command (1) Ship 031 Washington, D. C. 20360

Naval Ship Systems Command (1) Ship 035 Washington, D. C. 20360

U. S. Naval Weapons Laboratory (1) Dahlgren, Virginia 22448

Naval Electronic Systems Command (2) ELEX 03, Room 2046 Munitions Building Department of the Navy Washington, D. C. 20360

Head, Technical Services Division (1) Naval Investigative Service Headquarters 4420 North Fairfax Drive Arlington, Virginia 22203

Commander (2) U. S. Naval Ordnance Laboratory Attn: Librarian White Oak, Maryland 21502

Commanding Officer (2) Office of Naval Research Branch Office Box 39 FPO New York, New York 09510

Commanding Officer (1) Office of Naval Research Branch Office 219 South Dearborn Street Chicago, Illinois 60604

Commanding Officer (1) Office of Naval Research Branch Office 495 Summer Street Boston, Massachusetts 02210 Commander (1) Naval Air Development Center Johnsville, Warminster, Pa. 18974

Commanding Officer (1) Naval Training Device Center Orlando, Florida 32813

Commander (Code 753) (1) Naval Weapons Center Attn: Technical Library China Lake, California 93555

Commanding Officer (1) Naval Weapons Center Corona Laboratories Attn: Library Corona, California 91720

Commander (1) U. S. Naval Missile Center Point Mugu, California 93041

W. A. Eberspacher, Associate Head (1) Systems Integration Division Code 5340A, Box 15 U. S. Naval Missile Center Point Mugu, California 93041

Commander (2)
Naval Electronics Laboratory Center
Attn: Library
San Diego, California 92152

Deputy Director and Chief Scientist (1) Office of Naval Research Branch Office 1030 East Green Street Pasadena, California 91101

Library (Code 2124) (1) Technical Report Section Naval Postgraduate School Monterey, California 93940

Glen A. Myers (Code 52 Mv) (1) Assoc. Professor of Electrical Engineering Naval Postgraduate School Monterey, California 93940

Commanding Officer & Director (1) U.S. Naval Underwater Sound Laboratory Fort Trumbull New London, Connecticut 05840

Commanding Officer (1) Naval Avionics Facility Indianapolis, Indiana 46241

#### OTHER GOVERNMENT AGENCIES

Dr. H. Harrison, Code RRE (1) Chief Electrophysics Branch National Aeronautics and Space Administration Washington, D. C. 20546

NASA Lewis Research Center Attn: Library 21000 Brookpark Road Cleveland, Ohio 44135

Los Alamos Scientific Laboratory (1) Attn: Reports Library P.O. Box 1663 Los Alamos, New Mexico 87544

Federal Aviation Administration (1) Attn: Admin Stds Div (MS-110) 800 Independence Avenue S. W. Washington, D. C. 20590

Mr. M. Zane Thornton, Chief (1) Network Engineering, Communications and Operations Branch Lister Hill National Center for Blomedical Communications 8600 Rockville Pike Bethesda, Maryland 20014

U. S. Post Office Department (1) Library - Room 6012 12th & Pennsylvania Ave. N. W. Washington, D. C. 20260

#### NON-GOVERNMENT AGENCIES

Director (1)
Research Laboratory of Electronics
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139

Mr. Jerome Fox, Research Coordinator (1) Polytechnical Institute of Brooklyn 333 Jay Street Brooklyn, New York 11201 Director (1)
Columbia Radiation Laboratory
Columbia University
538 West 120th Street
New York, New York 10027

Director (1) Coordinated Science Laboratory University of Illinois Urbana, Illinois 61801

Director (1) Stanford Electronics Laboratories Stanford University Stanford, California 94305

Director (1) Microwave Physics Laboratory Stanford University Stanford, California 94305

Director (1) Electronics Research Laboratory University of California Berkeley, California 94720

Director (1) Electronic Sciences Laboratory University of Southern California Los Angeles, California 90007

Director (1) Electronics Research Center The University of Texas at Austin Austin, Texas 78712

Division of Engineering and Applied Physics 210 Pierce Hali Harvard University Cambridge, Massachusetts 02138

Dr. G. J. Murphy (1)
The Technological Institute
Northwestern University
Evanston, Illinois 60201

Dr. John C. Hancock, Head (1) School of Electrical Engineering Purdue University Lafayette, Indiana 47907

Dept. of Electrical Engineering (1) Texas Technological College Lubbock, Texas 79409

Aerospace Corporation
P. O. Box 95085
Los Angeles, California 90045
Attn: Library Acquisitions Group

Professor Nicholas George (1) California Inst. of Technology Pasadena, California 91109

Aeronautics Library (1) Graduate Aeronautical Laboratories California Institute of Technology 1201 E. California Blvd, Pasadena, California 91109

The Johns Hopkins University (1) Applied Physics Laboratory Attn: Document Librarian 8621 Georgia Avenue Silver Spring, Maryland 20910

Hunt Library (1) Carnegie-Mellon University Schenley Park Pittsburgh, Pennsylvania 15213

Dr. Leo Young (1) Stanford Research Institute Menlo Park, California 94025

School of Engineering Sciences (1) Arizona State University Tempe, Arizona 85281

Engineering & Mathematical Sciences Library (1) University of California at Los Angeles 405 Hilgred Avenue Los Angeles, California 90024

The Library (1)
Government Publications Section
University of California
Santa Barbara, California 93106

Carnegie Institute of Technology (1) Electrical Engineering Department Pittsburgh, Pennsylvania 15213

Professor Joseph E. Rowe (1) Chairman, Dept. of Electrical Engineering The University of Michigan Ann Arbor, Michigan 48104 New York University (1) College of Engineering New York, New York 10019

Syracuse University (1)
Dept. of Electrical Engineering
Syracuse, New York 13210

Yale University (1)
Engineering Department
New Haven, Connecticut 06520

Airborne Instruments Laboratory (1) Deerpark, New York 11729

Raytheon Company Attn: Librarian Bedford, Massachusetts 01730

Raytheon Company (1) Research Division Library 28 Seyon Street Waltham, Massachusetts 02154

Dr. Sheldon J. Welles (1)
Electronic Properties Information
Center
Mail Station E-175
Hughes Aircraft Company
Culver City, California 90230

Dr. Robert E. Fontana (1)
Systems Research Laboratories Inc.
7001 Indian Ripple Road
Dayton, Chio 45440

Nuclear Instrumentation Group (1) Bldg. 29, Room 101 Lawrence Radiation Laboratory University of California Berkeley, California 94720

Sylvania Electronic Systems (1) Applied Research Laboratory Attn: Documents Librarian 40 Sylvan Road Waitham, Massachusetts 02154

Hollander Associates (1) P. O. Box 2276 Fullerton, California 92633

Illinois Institute of Technology (1)
Department of Electrical Engineering
Chicago, Illinois 60616

The University of Arizona (1) Dept. of Electrical Engineering Tucson, Arizona 85721

Utah State University (1) Dept. of Electrical Engineering Logan, Utah 84321

Case Institute of Technology (1) Engineering Division University Circle Cieveland, Ohio 44106

Lincoln Laboratory (1)
Massachusetts Institute of
Technology
Lexington, Massachusetts 02173

The University of Iowa (1) The University Libraries Iowa City, Iowa 52240

Lenkurt Electric Co., Inc. (1) 1105 County Road San Carlos, California 94070 Attn: Mr. E. K. Peterson

Communications & Electronics Division Union Meeting and Jolly Roads Blue Bell, Pennsvylania 19422

Union Carbide Corporation (1) Electronic Division P.O. Box 1209 Mountain View, California 94041

Dept. of Electrical Engineering (1) Rice University Houston, Texas 77001

Research Laboratories for the Engineering Sciences (1) School of Engineering and Applied Science University of Virginia Charlottesville, Virginia 22903

Dept. of Electrical Engineering (1) College of Engineering and Technology Ohio University Athens, Ohio 45701 Security Classification

DOCUMENT CONTROL DATA - R&D (Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)							
1 ORIGINATING ACTIVITY (Corporate author) The University of Texas at Austin		2. REPORT SECURITY CLASSIFICATION UNCLASSIFIED					
							Electronics Research Center
Austin, Texas 78712	<u></u>						
3 REPORT TITLE							
THE STABILITY OF COUPLED RENEWAL-	DIFFERENTIAL E	QUATIO	ONS WITH				
ECONOMETRIC APPLICATIONS	• •	•	•				
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)							
Scientific Interim							
5. AUTHOR(S) (Last name, first name, initial)	• •						
Ronald P. Rhoten	·						
J. K. Aggarwal							
6 REPORT DATE	74. TOTAL NO. OF P	AGES	76. NO. OF REFS				
15 July 1969	61		19				
8a. CONTRACT OR GRANT NO.	9a. ORIGINATOR'S REPORT NUMBER(S)						
AF-AFOSR 67-0766E	JSEP, Tech	eport No. 69					
6. PROJECT NO. 4751							
47.01	AL OTHER REPORT	10(8) (4 av	other numbers that may be applicated				
6144501F	9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)						
d. 681305	AFOSR 69-2	2133TR					
10. A VAIL ABILITY/LIMITATION NOTICES	-						
1. This document has been approved for	or public releas	e and s	ale; its distribution				
is unlimited.							
11. SUPPLEMENTARY NOTES	12. SPONSORING MILL	TARY ACTI	VITY ISEP through				
THE SUFFE CHARTERY NOTES	12. SPONSORING MILITARY ACTIVITY JSEP through AF Office of Scientific Research (SREE)						
TECH, OTHER	CH, OTHER 1400 Wilson Boulevard						
	Arlington, Vi	'irginia 22209					
13. ABSTRACT							

This work presents concepts and results in the fields of mathematical modelling, economics and stability analysis. A coupled renewal-differential equation structure is presented as a modelling form for systems possessing hereditary characteristics, and this structure is applied to a model of the Austrian theory of business cycles. For realistic conditions, the system is shown to have an infinite number of poles, and conditions are presented which are both necessary and sufficient for all poles to lie strictly in the left half

plane.

Security Classification

14 KEY WORDS	LINK A		LINK B		LINK &		
KET WORDS		ROLE	WT	ROLE	WT	ROLE	WT
TIME DELAY							
BUSINESS CYCLES							-
RENEWAL-DIFFERENTIAL EQUATIONS	5						
TRANSCENDENTAL ZERO LOCATIONS	}						t
	·						
	·		İ				
	·						
				{			

#### INSTRUCTIONS

- 1. ORIGINATING ACTIVITY: Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (corporate author) issuing the report.
- 2a. REPORT SECURITY CLASSIFICATION: Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.
- 2b. GROUP: Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.
- 3. REPORT TITLE: Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.
- 4. DESCRIPTIVE NOTES: If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.
- 5. AUTHOR(S): Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.
- 6. REPORT DATE: Enter the date of the report as day, month, year, or month, year. If more than one date appears on the report, use date of publication.
- 7a. TOTAL NUMBER OF PAGES: The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.
- 7b. NUMBER OF REFERENCES: Enter the total number of references cited in the report.
- 8a. CONTRACT OR GRANT NUMBER: If appropriate, enter the applicable number of the contract or grant under which the report was written.
- 8b, 8c, & 8d. PROJECT NUMBER: Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.
- 9a. ORIGINATOR'S REPORT NUMBER(S): Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.
- 9b. OTHER REPORT NUMBER(\$): If the report has been assigned any other report numbers (either by the originator or by the sponsor), also enter this number(s).
- 10. AVAILABILITY/LIMITATION NOTICES: Enter any limitations on further dissemination of the report, other than those

imposed by security classification, using standard statements such as:

- "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known

- 11. SUPPLEMENTARY NOTES: Use for additional explanatory notes.
- 12. SPONSORING MILITARY ACTIVITY: Enter the name of the departmental project office or laboratory sponsoring (paying for) the research and development. Include address.
- 13. ABSTRACT: Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. KEY WORDS: Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, rules, and weights is optional